Ibuprofen dosage chart for infants and children 11 years and younger

Ibuprofen	If possible, use weight to dose; otherwise, use age.									
mL = milliliter mg = milligram	 <6 mos	12-17 lbs 6-11 mos	18-23 lbs 12-23 mos	24-35 lbs 2-3 yrs	36-47 lbs 4-5 yrs	48-59 lbs 6-8 yrs	60-71 lbs 9-10 yrs	72-95 lbs 11 yrs		
Infants' ibuprofen concentrated drops for example Concentrated Motrin° Infants' Drops 50 mg per 1.25 mL	Ask a healthcare professional	1.25 mL	1.875 mL	-	_	_	_	_		
Children's ibuprofen liquid for example Children's Advil* Suspension 100 mg per 5 mL	<	Ask a healthcard professional	€	5 mL	7.5 mL	10 mL	12.5 mL	15 mL		
Junior strength ibuprofen chewable tablets for example Junior Strength Advil* Chewables 100 mg per chewable tablet	<		Ask a healthcare professional		·	2 chewables	2½ chewables	3 chewables		

Important instructions

You may repeat the dose every 6 to 8 hours while symptoms last

Do **NOT** give more than 4 doses in 24 hours

Talk to your child's healthcare professional before giving any other product containing an NSAID*



*Nonsteroidal anti-inflammatory drug

- What you can do · Always read and follow the medicine label.
 - Find the dose for your child's current weight. If you are not sure of your child's weight, use age. As children grow, their doses change over time.
 - Use only the dosing device that came with the medicine (syringe or dosing cup).
 - · Wait the correct amount of time between doses. If you give another dose too soon, it could harm your child.

See acetaminophen dosage directions ▶



Get Relief Responsibly®

Acetaminophen dosage chart for infants and children 11 years and younger

Acetaminophen	If possible, use weight to dose; otherwise, use age.												
mL = milliliter mg = milligram	6-11 lbs 0-3 mos	12-17 lbs 4-11 mos	18-23 lbs 12-23 mos	24-35 lbs 2-3 yrs	36-47 lbs 4-5 yrs	48-59 lbs 6-8 yrs	60-71 lbs 9-10 yrs	72-95 lbs					
Infants' acetaminophen liquid for example Infants' Tylenol* Oral Suspension 160 mg per 5 mL		mL thcare professional ecommendation ex		5 mL	_	_	_	_		Important instructions You may repeat			
Children's acetaminophen liquid for example Children's Tylenol* Oral Suspension 160 mg per 5 mL	<u> </u>	Ask a healthcare professional	·	5 mL	7.5 mL	10 mL	12.5 mL	15 mL		the dose every 4 hours while symptoms last Do NOT give more than 5 doses in 24 hours			
Children's 80 mg acetaminophen tablets 80 mg per chewable tablet	<u></u>	Ask a healthcare professional		2 chewables	3 chewables	4 chewables	5 chewables	6 chewables		Do NOT use with any other product containing acetaminophen			
Children's or Junior 160 mg acetaminophen tablets for example Children's Tylenol* Chewables 160 mg per chewable tablet		Ask a healthcare professional		1 chewable	1½ chewables	2 chewables	2½ chewables	3 chewables					

What you can do • Always read and follow the medicine label.

- Find the dose for your child's current weight. If you are not sure of your child's weight, use age. As children grow, their doses change over time.
- Use only the dosing device that came with the medicine (syringe or dosing cup).
- Wait the correct amount of time between doses. If you give another dose too soon, it could harm your child.

See ibuprofen dosage directions ▶



Get Relief Responsibly®